


HOW WELL DOES QCD WORK FOR PHOTON-PHOTON COLLISIONS?

THORSTEN WENGLER

CERN, EP-division, 1211 Geneva 23, Switzerland



The performance of QCD in describing hadronic photon-photon collisions is investigated in the light of recent measurements from LEP on di-jet production, light hadron transverse momentum spectra, and heavy quark production.

1 Jet Production

Differential di-jet cross sections have been measured by DELPHI¹ and OPAL². DELPHI has analysed 220 pb⁻¹ of data taken at $\sqrt{s_{ee}}=192\text{--}202$ GeV using a cone algorithm with $R=1$ to define the jets. The first preliminary results of the di-jet cross sections as a function of the jet transverse energy and the jet pseudorapidity are found to be consistent with a previous measurement by OPAL³. OPAL has used 593 pb⁻¹ from $\sqrt{s_{ee}}=189$ GeV to 209 GeV to study the performance of NLO perturbative QCD and the hadronic structure of the photon in di-jet production. The jets are reconstructed using an inclusive $k_{\perp}$ clustering algorithm⁴. The two jets are used to estimate the fraction of the photon momentum participating in the hard interaction as $x_{\gamma}^{\pm} \equiv \sum_{\text{jets}=1,2} (E^{\text{jet}} \pm p_z^{\text{jet}}) / \sum_{\text{hfs}} (E \pm p_z)$ where p_z is the momentum component along the z axis of the detector and E is the energy of the jets or objects of the hadronic final state (hfs). In single (double) resolved processes, where one (both) of the photons interacts via a fluctuation into a hadronic state, the hard interaction is accompanied by one or two remnant jets, and one or both x_{γ} are smaller than one. Differential cross sections as a function of x_{γ} or in regions of x_{γ} are therefore a sensitive probe of the structure of the photon. The left plot in Figure 1 shows the differential di-jet cross section as a function of the mean transverse energy $\bar{E}_T^{\text{jet}}$ of the di-jet system for the full x_{γ} -range, for either x_{γ}^{+} or $x_{\gamma}^{-} < 1$ (dominated by single resolved processes), or $x_{\gamma}^{\pm} < 1$ (dominated by double resolved processes). The prediction of perturbative QCD in NLO⁵ using the GRV-GHO⁶ parton densities is compared to the data after hadronisation corrections have been applied to the calculation. The calculation is in good agreement with the data, except for being too low at small $\bar{E}_T^{\text{jet}}$ for $x_{\gamma}^{\pm} < 1$. In this region the contribution of the so




Figure 1: Differential di-jet cross sections as measured by OPAL.

called underlying event as described by the concept of multiple parton interactions (MIA) is expected to be largest. This contribution is not included in the NLO calculation. The three plots on the right hand side of Figure 1 show the differential cross section as a function of x_γ for the three regions in $x_\gamma^+-x_\gamma^-$ -space described above. The shaded histogram on the bottom of each of the three plots indicates the contribution of MIA to the cross section as obtained from the PYTHIA ⁷ MC generator. It is evident especially for $x_\gamma^\pm < 1$ that the MIA contribution is of about the same size as the discrepancy between the measurement and the NLO prediction. Furthermore it is interesting to observe that there is next to no MIA contribution to the cross section if either x_γ^+ or x_γ^- is required to be less than one, while the sensitivity to the photon structure at small x_γ is retained. As one would expect also the agreement of the NLO calculation with the measurement is best in this case. With these measurements one is therefore able to disentangle the hard subprocess from soft contributions and make the firm statement that NLO perturbative QCD is adequate to describe di-jet production in photon-photon collisions. At the same time a different sub-set of observables can be used to study in more detail the nature of the soft processes leading to the underlying event.

2 Transverse Momentum Spectra of Light Hadrons

The L3 collaboration has recently presented new results on the inclusive production of K_S^0 and neutral and charged pions in 414 pb^{-1} of data taken at $\sqrt{s_{\text{ee}}}$ from 189 GeV to 202 GeV ⁸. Compared to the jet measurements described above, inclusive hadron production offers a complementary signature to study the dynamics in hadronic photon-photon collisions. Predictions are available in NLO perturbative QCD which make use of fragmentation functions to translate the partonic cross sections to the observables measured. The uncertainty associated with the hadronisation correction commonly applied to the calculation of partonic NLO jet cross sections can hence be avoided, but is of course replaced by any uncertainty attached to the determination of the fragmentation functions. The new measurements can be shown to be consistent with results obtained previously by the OPAL collaboration ⁹ where both measurements overlap in phase space. The new L3 data significantly extends the older measurements towards high transverse momenta for the production of charged and neutral pions. As can be seen in Figure 2 it is




Figure 2: The differential cross section as a function of the transverse momentum of the hadron for neutral pions as measured by L3 (left) and the total cross section of charm production (ALEPH, DEPHI, L3, OPAL) and beauty production (L3, OPAL) in photon-photon collisions (right).

in this region that a significant discrepancy between the measurement and the QCD prediction occurs. The left plot in Figure 2 shows the differential cross section for inclusive π^0 -production as a function of the transverse momentum p_T of the π^0 . For $p_T > 8$ GeV the data exhibits a significantly harder spectrum than predicted by NLO QCD. This behaviour is confirmed by the independent measurement of charged pions. This discrepancy is particularly remarkable as it occurs at high transverse momenta of the produced hadrons, which in turn suggests an underlying high momentum partonic process. Under these conditions one expects the perturbative calculation to be reliable due to the relatively small value of the strong coupling constant. The discrepancy also appears to be in contradiction with the good agreement of the NLO QCD calculation obtained for the jet measurements described above, which should be sensitive to a similar set of processes and mainly differs in that fragmentation functions are not needed here. A possible direction for a further investigation of this anomaly may be derived from the OPAL measurement⁹. Here the data is presented in four regions of the invariant hadronic mass W . For $W < 50$ GeV a similar tendency of a harder p_T spectrum in the data then in the theory can be observed. For $W > 50$ GeV, however, the agreement is good. It would therefore be interesting to repeat these measurements for the full LEP2 data set in several regions of W .

3 Heavy Flavour Production

Yet another test of the performance of perturbative QCD in photon-photon collisions is the production of heavy quarks. Due to the large physical scale set by the charm or beauty mass one can expect the theoretical prediction from perturbative QCD to be reliable. A significant part of the cross section at LEP2 energies is predicted to be from resolved processes, in particular the photon-gluon fusion $\gamma g \rightarrow c\bar{c}(b\bar{b})$. The production rate of charm and beauty quarks therefore depends not only on their mass but also on the gluon density in the photon. The latest results from the LEP collaborations on charm production use identified charged D^* meson for charm tagging.

L3¹⁰ has analysed 683 pb⁻¹ at $\sqrt{s_{ee}}$ =183-209 GeV, DELPHI¹¹ 458 pb⁻¹ at $\sqrt{s_{ee}}$ =183-209 GeV, OPAL¹² 428 pb⁻¹ at $\sqrt{s_{ee}}$ =183-202 GeV, and ALEPH¹³ 236 pb⁻¹ at $\sqrt{s_{ee}}$ =183-189 GeV. The smallest uncertainties are to be expected if one compares the measurements and the theoretical predictions for the restricted phase space corresponding to the experimental acceptances, as any extrapolation to the total charm production cross section introduces additional assumptions. The differential cross section of charged D* production as a function of the D* transverse momentum is in good agreement among the four experiments, with the possible exception of a slightly harder spectrum for ALEPH. NLO perturbative QCD using massive quarks¹⁴ and a charm mass of m_c =1.5 GeV describes the data well. This translates to similar agreement of the total charm production cross section with the QCD prediction¹⁴ as can be seen in the right plot of Figure 2. In the same figure the results are shown for the total production cross section of $b\bar{b}$ as obtained by the L3 and the OPAL collaborations. Both measurements are based on the higher transverse momentum of the lepton in semileptonic b-decays as compared to those of lighter quarks. L3 has analysed 410 pb⁻¹ at $\sqrt{s_{ee}}$ =189-202 GeV using both the electron and the muon signatures¹⁵, OPAL has analysed 371 pb⁻¹ at $\sqrt{s_{ee}}$ =189-202 GeV using the muon channel¹⁶. The plot demonstrates that there is good agreement between the experimental results. The prediction of perturbative QCD in NLO however significantly underestimates the measurements. Similar deficiencies have been observed in b-production in $p\bar{p}$ - and ep-collisions. A recent analysis of the b-fragmentation functions used in these calculations suggests that much of this discrepancy can be recovered at least for $p\bar{p}$ -collisions¹⁷. This analysis considers also the higher moments of the differential distributions of the average b-quark energy fraction carried by the B-meson. Using recent LEP and SLD data in this way to fix the b-fragmentation the agreement of the perturbative calculation with the $p\bar{p}$ -data becomes acceptable. One might hope that a similar technique applied to ep- and photon-photon-collisions will also improve the agreement of the theory with the measurements.

References

1. J. Mašik (DELPHI coll.), Proc. of PHOTON 2001, Ascona, World Scientific.
2. T. Wengler (OPAL coll.), Proc. of PHOTON 2001, Ascona, World Scientific.
3. OPAL Coll., G. Abbiendi *et al.*, *Euro. Phys. Journal C* **10**, 547 (1999).
4. S. Catani, Yu.L. Dokshitzer, M.H. Seymour, B.R. Webber, *Nucl. Phys. B* **406**, 187 (1993); S.D. Ellis, D.E. Soper, *Phys. Rev. D* **48**, 3160 (1993).
5. M. Klasen, T. Kleinwort, G. Kramer, *Euro. Phys. J. Direct C* **1**, 1 (1998); M. Klasen, private communication.
6. M. Glück, E. Reya, A. Vogt, *Phys. Rev. D* **45**, 3986 (1992); M. Glück, E. Reya, A. Vogt, *Phys. Rev. D* **46**, 1973 (1992).
7. T. Sjöstrand, *Comp. Phys. Comm.* **82**, 74 (1994); T. Sjöstrand, LUND University Report, LU-TP-95-20 (1995).
8. L3 Coll., P. Achard *et al.*, *Phys. Lett. B* **524**, 44 (2002); L3 note 2730, 2002.
9. OPAL Coll., K. Ackerstaff *et al.*, *Euro. Phys. Journal C* **6**, 253 (1999).
10. L3 Coll., P. Achard *et al.*, CERN-EP-2002-012, submitted to *Phys. Lett. B*.
11. M. Chapkin, V. Obraztsov, A. Sokolov (DELPHI Coll.), Proc. of PHOTON 2000, AIP conf. proc., v. 571, 252 (2000).
12. OPAL Coll., G. Abbiendi *et al.*, *Euro. Phys. Journal C* **16**, 579 (2000).
13. ALEPH Coll., D. Buskulic *et al.*, *Phys. Lett. B* **355**, 595 (1995); ALEPH note 031, 2000.
14. V. Andreev (L3 coll.), Proc. of PHOTON 2001, Ascona, World Scientific.
15. L3 Coll., M. Acciari *et al.*, *Phys. Lett. B* **503**, 10 (2001).
16. Á. Csilling, (OPAL Coll.), Proc. of PHOTON 2000, AIP conf. proc., v. 571, 276 (2000).
17. M. Cacciari, these proceedings.